arXiv:1303.5283v1 [physics.gen-ph] 19 Mar 2013

On the electron to proton mass ratio

Ole L. Trinhammer
Department of Physics, Technical University of Denmark,
Fysikvej, Building 307,
DK-2800 Kongens Lyngby, Denmark.

We derive an expression for the electron to nucleon mass ratio from a reinterpreted Kogut-Susskind Hamiltonian to describe interior baryon dynamics. We use the classical electron radius as our fundamental length scale. Based on expansions on trigonometric Slater determinants for a neutral state a specific numerical result is found to be less than three percent off the experimental value for the neutron.

1 The result and the model

The ratio we get between the electron mass and the proton mass is

$$\frac{m_e}{m_p} = \frac{\alpha}{\pi}\frac{1}{\mathrm{E}}, \tag{1}$$

where α is the fine structure constant and $\mathrm{E} = E/\Lambda$ is the dimensionless ground state eigenvalue of a reinterpreted Kogut-Susskind Hamiltonian [1]

$$\frac{\hbar c}{a}\left[-\frac{1}{2}\Delta + \frac{1}{2}\mathrm{Tr}\chi^2\right]\Psi(u) = E\Psi(u). \tag{2}$$

with Manton's [2] representation independent [3] action for a link variable [4] used now as a potential. The energy scale $\Lambda \equiv \hbar c/a$ corresponds to a fundamental length scale a, which we shall relate to the classical electron radius [5]. It is not a new idea to suggest the classical electron radius as a fundamental length in elementary particle physics [6, 7]. We assume (2) to describe the baryon spectrum with configuration variable $u = e^{i\chi}$ in the Lie group u(3) and identify the ground state with the proton. We have derived an operable expression for the Laplacian Δ in [8]. We have presented our model (2) in brief form in [9] and worked through the details to compare with the relative neutron to proton mass shift, baryon resonance spectroscopy and parton distribution functions in [10]. Our configuration space is "orthogonal" to the laboratory space wherefore (2) describes a truly interior dynamics which may be projected on laboratory space parameters through the eigenangles θ_j parametrizing the eigenvalues $e^{i\theta_j}$ of u. The projection introduces the dimensionful scale a, thus

$$x_j = a\theta_j. \tag{3}$$

Conjugate to the space (angle) parameters in (3) are canonical momentum (interior action) operators

$$p_j = -i\hbar\frac{1}{a}\frac{\partial}{\partial\theta_j} = \frac{\hbar}{a}T_j. \tag{4}$$

The toroidal generators T_j induce coordinate fields ∂_j according to the above mentioned eigenangle parametrization of the u(3) torus. In general the nine generators T_k of u(3) namely induce coordinate fields as follows



FIG. 1: Projection from configuration space to laboratory space.

$$\partial_k = \frac{\partial}{\partial\theta}ue^{i\theta T_k}|_{\theta=0} = uiT_k. \tag{5}$$

The remaining six off-toroidal generators are important in the baryon spectroscopy phenomenology resulting from (2) since they take care of spin and flavour degrees of freedom. They need not be specified further in the present context than to mention that they can be represented by the six off-diagonal Gell-Mann generators and lead to a "centrifugal" potential in the polar decomposition of the Laplacian given in [8] and used in [9, 10].

2 The derivation

We us assume the relativistic relation between energy E, momentum p and rest mass m to hold in general, thus

$$E^2 = p^2c^2 + m^2c^4, \tag{6}$$

Now consider a baryon at rest. We imply two complementary views: Pure energy interior space with $m = 0$ gives an *allospatial* energy relation

$$E_a^2 = p_a^2c^2 \tag{7}$$

and complete rest with $p = 0$ in laboratory space gives the rest energy

$$E_0^2 = m^2c^4. \tag{8}$$

Both views relate to the same eigenvalue E of our Schrödinger equation and thus

$$p_a c = mc^2. \tag{9}$$

The quantization inherent in the concept of the de Broglie wavelength then gives at rest

$$\lambda_0 = \frac{h}{p_a} = 2\pi\frac{\hbar}{mc} = 2\pi\frac{\hbar c}{\mathrm{E}\Lambda} = \frac{2\pi a}{\mathrm{E}}. \tag{10}$$

The third expression is the Compton wavelength and the last expression indicates an orbital mapping from interior space to laboratory space. From the well known relation for the fine structure constant [11]

$$\alpha = \frac{1}{4\pi\epsilon_0}\frac{e^2}{\hbar c} \tag{11}$$

and the classical electron radius [5]

$$\frac{1}{4\pi\epsilon_0}\frac{e^2}{R_0} = m_ec^2 \tag{12}$$

we get

$$R_0 = \frac{\alpha\hbar c}{m_ec^2}. \tag{13}$$

Now a shortest geodesic to track along the full extension of the u(3) maximal torus runs from origo at $u = I$ where all eigenvalues are $1 = e^{i0}$ to $u = -I$ where all eigenvalues are $-1 = e^{i\pi}$, see also fig. 1. When for instance the neutron decays to the proton - and the electron is created as a "peel off" - the topological change in the interior baryon state maps by projection to laboratory space. From (3) we then have

$$R_0 = a\pi. \tag{14}$$

With (14) and (13) in (10) we readily find (1).

3 Bases for solutions

In [9, 10] we report on exact calculations for neutral states where the toroidal part of the wavefunction is expanded on Slater determinants

$$b_{pqr} = \epsilon_{ijk}\cos p\theta_i \sin q\theta_j \cos r\theta_k \tag{15}$$

with suitably chosen integers p, q, r. The result is $\mathrm{E} = 4.38$ for a neutral ground state m_0 corresponding to $m_e/m_0 = 1/1885$ which is within 3 percent of the value $m_e/m_n = 1/1838.6...$ based on experimental data for the electron and the neutron in [12]. Work is in progress to solve (2) for the charged proton by antisymmetrized parity definite expansions on Bloch states [13]

$$g_{pqr} = e^{i\boldsymbol{\kappa}\cdot\boldsymbol{\theta}}e^{ip\theta_1}e^{iq\theta_2}e^{ir\theta_3}, \tag{16}$$

where $\boldsymbol{\kappa} = (\kappa_1, \kappa_2, \kappa_3)$ are appropriately chosen Bloch vectors and $\boldsymbol{\theta} = (\theta_1, \theta_2, \theta_3)$.

4 Comments

It is not clear whether the minor but still significant discrepancy in our value for m_e/m_n is due to indefinite parity of the base set (15) or due to corrections needed in (14). It is somewhat surprising that when applied to a neutral state expression (1) gives a result just three percent off the experimental value for the electron to neutron value. Surprising since the classical electron radius on which we base our prediction is normally presented to be just an order of magnitude scale for strong interaction phenomena [14]. It should be noted though that in (3) we have introduced a well defined projection to space parameters. However, at the present level of understanding it seems wise to quote L. D. Landau and E. M. Lifshitz: "...it is impossible within the framework of classical electrodynamics to pose the question whether the total mass of the electron is electrodynamic." [5]

Conclusion

We have derived an expression for the electron to nucleon mass ratio from a reinterpreted lattice gauge theory Hamiltonian. A specific calculation for the neutron from expansions on Slater determinants of indefinite parity gives a result less than three percent off the experimental value. The proximity of prediction to observation should encourage further study within the framework of the Hamiltonian model presented. Work is in progress to establish a complete Bloch wave base for expansions with suitable symmetry requirements implied by the potential and the antisymmetry under interchange of colour degrees of freedom.

Acknowledgment

I am grateful to the late Victor F. Weisskopf for posing the inspiring problem of m_e/m_p to us summer students at CERN in Geneva 1976.

References

[1] J. B. Kogut and L. Susskind, *Hamiltonian formulation of Wilson's lattice gauge theories*, Phys. Rev. **D11** (2), 395, (1975).

[2] N. S. Manton, *An Alternative Action for Lattice Gauge Theories*, Phys. Lett. **B96**, 328-330 (1980).

[3] J. Milnor, Morse Theory, Ann. of Math. Stud. 51, 1 (1963).

[4] O. Trinhammer, *Infinite N phase transitions in one-plaquette (2+1)-dimensional models of lattice gauge theory with Manton's action*, Phys. Lett. **B129** (3, 4), 234-238 (1983).

[5] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields, Course of Theoretical Physics* Vol. 2, 4th ed, (Elsevier Butterworth-Heinemann, Oxford 2005), p.97.

[6] W. Heisenberg, *Über die in der Theorie der Elementarteilchen auftretende universelle Länge*, Ann. d. Phys. (5) 32, 20-33. (1938).

[7] W. Heisenberg discussed in J. Mehra and H. Rechenberg, *The Historical Development of Quantum Theory - The Conceptual Completion and the Extensions of Quantum Mechanics 1932-1941*, Vol. 6 (2), (Springer, New York 2001). p.956.

[8] O. L. Trinhammer and G. Olafsson, *The Full Laplace-Beltrami operator on U(N) and SU(N)*, arXiv:9901002v2 [math-ph] (1999/2012).

[9] O. L. Trinhammer, *Neutron to proton mass difference, parton distribution functions and baryon resonances from a simple abstract model*, arXiv:1302.1779 [hep-th] (2013).

[10] O. L. Trinhammer, *Neutron to proton mass difference, parton distribution functions and baryon resonances from dynamics on the Lie group u(3)*, arXiv:1109.4732v3 [hep-th] (2011/2012).

[11] J. J. Sakurai, *Advanced Quantum Mechanics*, 10th printing, (Addison-Wesley, United Kingdom 1967/1984), p.12.

[12] J. Beringer et al. (Particle Data Group), *Review of Particle Physics*, Phys. Rev. **D86**, 010001 (2012), p.30, p.79.

[13] Karsten W. Jacobsen, *private communication*, Technical University of Denmark, June 2012.

[14] W. Heisenberg, op. cit. p.20.